arXiv:1401.1296v1 [math.PR] 7 Jan 2014

Asymptotics of the probability distributions of the first hitting times of Bessel processes

Yuji Hamana[a] and Hiroyuki Matsumoto[b] [1]

[a]Department of Mathematics, Kumamoto University, Kurokami 2-39-1, Kumamoto 860-8555, Japan
[b]Department of Physics and Mathematics, Aoyama Gakuin University, Fuchinobe 5-10-1, Sagamihara 252-5258, Japan

Abstract. The asymptotic behavior of the tail probabilities for the first hitting times of the Bessel process with arbitrary index is shown without using the explicit expressions for the distribution function obtained in the authors' previous works.

2010 *Mathematics Subject Classification*: 60G40
keywords: Bessel process, hitting time, tail probability

1. Introduction and main results

Let $\mathbb{P}_a^{(\nu)}$ be the probability law on the path space $W = C([0,\infty);\mathbf{R})$ of a Bessel process with index $\nu \in \mathbf{R}$ or dimension $\delta = 2(\nu+1)$ starting from $a > 0$. For $w \in W$ we denote the first hitting time to $b > 0$ by $\tau_b = \tau_b(w)$:

$$\tau_b = \inf\{t > 0; w(t) = b\}.$$

In recent works [4, 5] the authors have shown explicit forms of the distribution function and the density of the distribution of τ_b under $\mathbb{P}_a^{(\nu)}$ in the case where $0 < b < a$. The other case, which is easier since we do not need to consider a natural boundary, has been known by Kent [6]. See also Gettor-Sharpe [2].

When $\nu > 0$ and $b < a$, it is shown in [5] that there exists a positive constant $C(\nu)$ such that

$$\mathbb{P}_a^{(\nu)}(\tau_b > t) = 1 - \left(\frac{b}{a}\right)^{2\nu} + C(\nu)t^{-\nu} + o(t^{-\nu}).$$

[1]Corresponding author.
E-mail adress: matsu(at)gem.aoyama.ac.jp(H.Matsumoto)

The constant $C(\nu)$ may be expressed explicitly and we could treat all the cases. However, we need to consider separately the case where δ is an odd integer and the expression for $C(\nu)$ is different from the othere cases. This is because the expression for the distribution function itself is different.

The aim of this note is to show that the constant $C(\nu)$ has the same simple expression also when δ is an odd integer by considering the asymptotics without using the explicit expressions for the distribution functions obtained in [5].

When $a < b$ and $\nu > 0$, the explicit expressions for $\mathbb{P}_a^{(\nu)}(\tau_b > t)$ has been shown in [6] and, from his result, it is easily shown that the tail probability decays exponentially. Hence we concentrate on the case of $b < a$.

Theorem 1. *Let $\nu > 0$ and $0 < b < a$. Then, as $t \to \infty$, it holds that*

$$\mathbb{P}_a^{(\nu)}(t < \tau_b < \infty) = b^{2\nu}\Big\{1 - \Big(\frac{b}{a}\Big)^{2\nu}\Big\}\frac{1}{\Gamma(1+\nu)(2t)^{\nu}} + O(t^{-\nu-\varepsilon})$$

for any $\varepsilon \in (0, \frac{\nu}{1+\nu})$, where Γ denotes the usual Gamma function.

Theorem 2. *Let $\nu < 0$ and $0 < b < a$. Then, as $t \to \infty$, it holds that*

$$\mathbb{P}_a^{(\nu)}(\tau_b > t) = a^{2|\nu|}\Big\{1 - \Big(\frac{b}{a}\Big)^{2|\nu|}\Big\}\frac{1}{\Gamma(1+|\nu|)(2t)^{|\nu|}} + O(t^{-|\nu|-\varepsilon})$$

for any $\varepsilon \in (0, \frac{\nu}{1+\nu})$.

When $\nu = 0$, it is known that

$$\mathbb{P}_a^{(0)}(\tau_b > t) = \frac{2\log(a/b)}{\log t} + o((\log t)^{-1}).$$

This identity has been discussed in [5] and we omit the details.

2. Proof of Theorem1

We assume $\nu > 0$ in this section. At first we give some lemmas. The first one is shown by Byczkowski and Ryznar [1].

Lemma 3. *There exists a constant C such that*

$$\mathbb{P}_a^{(\nu)}(t < \tau_b < \infty) \leqq Ct^{-\nu}.$$

Lemma 4. *If $0 < b < a$, one has*

$$\mathbb{P}_a^{(\nu)}(\tau_b > t) = 1 - \Big(\frac{b}{a}\Big)^{2\nu} + \mathbb{E}_a^{(\nu)}\Big[\Big(\frac{b}{R_t}\Big)^{2\nu}\mathbf{1}_{\{\inf_{0\leqq s\leqq t} R_s > b\}}\Big] \tag{1}$$

for any $t > 0$, where $\mathbb{E}_a^{(\nu)}$ is the excpectation with respect to $\mathbb{P}_a^{(\nu)}$ and $\{R_s\}_{s\geqq 0}$ denotes the coordinate process.

Proof. It is well known that

$$\mathbb{P}_a^{(\nu)}(\tau_b = \infty) = \mathbb{P}_a^{(\nu)}(\inf_{s\geqq 0} R_s > b) = 1 - \Big(\frac{b}{a}\Big)^{2\nu}.$$

By the Markov property of Bessel processes, we have for $t > 0$

$$\begin{aligned}\mathbb{P}_a^{(\nu)}(\tau_b = \infty) &= \mathbb{P}_a^{(\nu)}(\inf_{0\leqq s\leqq t} R_s > b \text{ and } \inf_{s\geqq t} R_s > b)\\ &= \mathbb{E}_a^{(\nu)}[\mathbb{P}_{R_t}^{(\nu)}(\tau_b = \infty)\mathbf{1}_{\{\inf_{0\leqq s\leqq t} R_s > b\}}]\\ &= \mathbb{E}_a^{(\nu)}\Big[\Big\{1 - \Big(\frac{b}{R_t}\Big)^{2\nu}\Big\}\mathbf{1}_{\{\inf_{0\leqq s\leqq t} R_s > b\}}\Big],\end{aligned}$$

which implies (1).

Lemma 5. *For any $a > 0$ and p with $0 < p < 1 + \nu$, it holds that*

$$\frac{\Gamma(1+\nu-p)}{\Gamma(1+\nu)}\frac{1}{(2t)^p}e^{-\frac{a^2}{2t}} \leqq \mathbb{E}_a^{(\nu)}[(R_t)^{-2p}] \leqq \frac{\Gamma(1+\nu-p)}{\Gamma(1+\nu)}\frac{1}{(2t)^p} + Ct^{-1-p} \qquad (2)$$

for $t \geqq 1$, where C is a positive constant independent of t.

Proof. By the explicit expression for the transition density of the Bessel process, we have

$$\mathbb{E}_a^{(\nu)}[(R_t)^{-2p}] = \int_0^\infty y^{-2p}\frac{1}{t}\Big(\frac{y}{a}\Big)^\nu y e^{-\frac{a^2+y^2}{2t}} I_\nu\Big(\frac{ay}{t}\Big)\mathrm{d}y,$$

where I_ν is the modified Bessel function of the first kind with index ν (cf [8]) given by

$$I_\nu(z) = \Big(\frac{z}{2}\Big)^\nu \sum_{n=0}^\infty \frac{(z/2)^{2n}}{\Gamma(n+1)\Gamma(1+\nu+n)} \quad (z \in \mathbf{R}\setminus(-\infty, 0)).$$

Hence, it is easy to get

$$\mathbb{E}_a^{(\nu)}[(R_t)^{-2p}] = \frac{1}{(2t)^p}e^{-a^2/2t}\sum_{n=0}^\infty \frac{a^{2n}\Gamma(n+\nu+1-p)}{\Gamma(n+1)\Gamma(1+n+\nu)(2t)^n}$$

and the assertion of the lemma.

Remark 6. The moments of R_t for fixed t have explicit expressions by means of the Whittaker functions (cf. [3], p.709), but it does not seem useful.

We are now in a position to give a complete proof of Theorem 1.

Proof of Theorem 1. By Lemma 4 we have

$$\mathbb{P}_a^{(\nu)}(t < \tau_b < \infty) = b^{2\nu}\mathbb{E}_a^{(\nu)}[(R_t)^{-2\nu}] - b^{2\nu}\mathbb{E}_a^{(\nu)}[(R_t)^{-2\nu}\mathbf{1}_{\{\tau_b\leqq t\}}].$$

For the first term we have shown in Lemma 5

$$\mathbb{E}_a^{(\nu)}[(R_t)^{-2\nu}] = \frac{1}{\Gamma(1+\nu)(2t)^\nu}(1 + O(t^{-1})).$$

Hence, if we could show

$$\mathbb{E}_a^{(\nu)}[(R_t)^{-2\nu}\mathbf{1}_{\{\tau_b\leqq t\}}] = \frac{1}{\Gamma(\nu+1)(2t)^\nu}\Big(\frac{b}{a}\Big)^{2\nu} + O\Big(\frac{1}{t^{\nu+\varepsilon}}\Big) \qquad (3)$$

for any $\varepsilon \in (0, \frac{\nu}{\nu+1})$, we obtain the assertion of the theorem.

For this purpose, we let $\alpha \in (0, \frac{1}{\nu+1})$, choose p satisfying

$$\frac{1}{1-\alpha} < p < \frac{1+\nu}{\nu}$$

and let q be such that $p^{-1} + q^{-1} = 1$. We devide the expectation on the right hand side of (3) into the sum of

$$I_1 = \mathbb{E}_a^{(\nu)}[(R_t)^{-2\nu}\mathbf{1}_{\{\tau_b \leqq t^{\alpha q}\}}] \quad \text{and} \quad I_2 = \mathbb{E}_a^{(\nu)}[(R_t)^{-2\nu}\mathbf{1}_{\{t^{\alpha q} < \tau_b \leqq t\}}]$$

We simply apply the Hölder inequality to I_2. Then we get

$$\begin{aligned} I_2 &\leqq \mathbb{E}_a^{(\nu)}[(R_t)^{-2\nu}\mathbf{1}_{\{t^{\alpha q} < \tau_b < \infty\}}] \\ &\leqq \left\{\mathbb{E}_a^{(\nu)}[(R_t)^{-2\nu p}]\right\}^{1/p}\left\{\mathbb{P}_a^{(\nu)}(t^{\alpha q} < \tau_b < \infty)\right\}^{1/q}. \end{aligned}$$

and, by Lemmas 3 and 5, we see that there exists a constant C_1 such that

$$I_2 \leqq C_1 t^{-\nu-\alpha\nu}.$$

In the following we denote by C_i's the constants independent of t.

For I_1, the strong Markov property of Bessel processes implies

$$I_1 = \int_0^{t^{\alpha q}} \mathbb{E}_b^{(\nu)}[(R_{t-s})^{-2\nu}]\mathbb{P}_a^{(\nu)}(\tau_b \in ds) = I_{11} + I_{12},$$

where

$$I_{11} = \int_0^{t^{\alpha q}} \frac{1}{2^\nu \Gamma(\nu+1)} \frac{1}{(t-s)^\nu} \mathbb{P}_a^{(\nu)}(\tau_b \in ds).$$

Since $\alpha q < 1$, Lemma 3 imples

$$|I_{12}| = |I_1 - I_{11}| \leqq \int_0^{t^{\alpha q}} \frac{C_2}{(t-s)^{\nu+1}} \mathbb{P}_a^{(\nu)}(\tau_b \in ds) \leqq \frac{C_3}{t^{\nu+1}}.$$

We devide I_{11} into the sum of

$$J_1 = \int_{t^\alpha}^{t^{\alpha q}} \frac{1}{2^\nu \Gamma(\nu+1)} \frac{1}{(t-s)^\nu} \mathbb{P}_a^{(\nu)}(\tau_b \in ds)$$

and

$$J_2 = \int_0^{t^\alpha} \frac{1}{2^\nu \Gamma(\nu+1)} \frac{1}{(t-s)^\nu} \mathbb{P}_a^{(\nu)}(\tau_b \in ds).$$

For J_1 we have by Lemma 3

$$0 \leqq J_1 \leqq \frac{C_4}{(t-t^{\alpha q})^\nu} \mathbb{P}_a^{(\nu)}(t^\alpha < \tau_b < \infty) \leqq \frac{C_5}{t^{\nu+\alpha\nu}}.$$

For J_2 we have

$$\begin{aligned} J_2 &\leqq \frac{1}{2^\nu \Gamma(\nu+1)(t-t^\alpha)^\nu} \mathbb{P}_a^{(\nu)}(\tau_b \leqq t^\alpha) \\ &\leqq \frac{1}{\Gamma(\nu+1)(2t)^\nu} \mathbb{P}_a^{(\nu)}(\tau_b < \infty) \frac{1}{(1-t^{-(1-\alpha)})^\nu} \\ &\leqq \frac{1}{\Gamma(\nu+1)(2t)^\nu} \Big(\frac{b}{a}\Big)^{2\nu} \Big(1 + \frac{C_6}{t^{1-\alpha}}\Big). \end{aligned}$$

On the other hand we have by Lemma 3

$$\begin{aligned} J_2 &\geqq \frac{1}{\Gamma(\nu+1)(2t)^\nu} \mathbb{P}_a^{(\nu)}(\tau_b \leqq t^\alpha) \\ &= \frac{1}{\Gamma(\nu+1)(2t)^\nu} \Big\{ \mathbb{P}_a^{(\nu)}(\tau_b < \infty) - \mathbb{P}_a^{(\nu)}(t^\alpha \leqq \tau_b < \infty) \Big\} \\ &\geqq \frac{1}{\Gamma(\nu+1)(2t)^\nu} \Big(\frac{b}{a}\Big)^{2\nu} - \frac{C_7}{t^\nu t^{\alpha\nu}}. \end{aligned}$$

Combining the above estimates, we obtain

$$\mathbb{E}_a^{(\nu)}[(R_t)^{-2\nu} \mathbf{1}_{\{\tau_b \leqq t\}}] = \frac{1}{\Gamma(\nu+1)(2t)^\nu} \Big(\frac{b}{a}\Big)^{2\nu} + \frac{1}{t^\nu} O\Big(\frac{1}{t^{\alpha\nu}} + \frac{1}{t} + \frac{1}{t^{1-\alpha}}\Big).$$

Since

$$0 < \alpha\nu < \frac{\nu}{\nu+1} < 1-\alpha < 1$$

and we can choose arbitrary α satifying this condition,

$$\mathbb{E}_a^{(\nu)}[(R_t)^{-2\nu} \mathbf{1}_{\{\tau_b \leqq t\}}] = \frac{1}{\Gamma(\nu+1)(2t)^\nu} \Big(\frac{b}{a}\Big)^{2\nu} + O\Big(\frac{1}{t^{\nu+\varepsilon}}\Big)$$

holds for any $\varepsilon \in (0, \frac{\nu}{\nu+1})$.

Now we have shown (3) and the assertion of Theorem 1.

3. Proof of Theorem 2

Theorem 2 is easily obtained from Theorem 1.

We recall explcit expressions for the Laplace transforms of the distributions of τ_b: for $\nu \in \mathbf{R}$, it is known ([2, 5]) that

$$\mathbb{E}_a^{(\nu)}[e^{-\lambda\tau_b}] = \Big(\frac{b}{a}\Big)^\nu \frac{K_\nu(a\sqrt{2\lambda})}{K_\nu(b\sqrt{2\lambda})}, \qquad \lambda > 0,$$

where K_ν is the modified Bessel function of the second kind. From this identity we easily obtain for $\nu > 0$

$$\mathbb{P}_a^{(-\nu)}(\tau_b \in dt) = \Big(\frac{a}{b}\Big)^{2\nu} \mathbb{P}_a^{(\nu)}(\tau_b \in dt).$$

Hence we get from Theorem 1

$$\begin{aligned}\mathbb{P}_a^{(-\nu)}(\tau_b > t) &= \left(\frac{a}{b}\right)^{2\nu} \mathbb{P}_a^{(\nu)}(t < \tau_b < \infty) \\ &= a^{2\nu}\Big\{1 - \left(\frac{b}{a}\right)^{2\nu}\Big\}\frac{1}{\Gamma(1+\nu)(2t)^\nu}(1 + o(1)).\end{aligned}$$

Acknowledgement. We thank Professor Yuu Hariya for valuable discussions.

References

[1] T. Byczkowski and M. Ryznar, Hitting distribution of geometric Brownian motion, Studia Math., **173** (2006), 19–38.

[2] R. K. Getoor and M. J. Sharpe, Excursions of Brownian motion and Bessel processes, Z. Wahr. Ver. Gebiete, **47** (1979), 83–106.

[3] I. S. Gradshteyn and I. M. Ryzhik, Table of Integrals, Series, and Products, 7th ed. (Academic Press, Amsterdam 2007).

[4] Y. Hamana and H. Matsumoto, The probability densities of the first hitting times of Bessel processes, J. Math-for-Industry, **4** (2012), 91–95.

[5] Y. Hamana and H. Matsumoto, The probability distributions of the first hitting times of Bessel processes, Trans. AMS, **365** (2013), 5237–5257.

[6] J. T. Kent, Eigenvalue expansions for diffusion hitting times, Z. Wahr. Ver. Gebiete, **52** (1980), 309–319.

[7] D. Revuz and M. Yor, *Continuous Martingales and Brownian Motion*, 3rd ed., Spronger-Verlag, 1999.

[8] G. N. Watson, *A Treatise on the Theory of Bessel Functions*, Reprinted of 2nd ed., Cambridge University Press, 1995.